

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

REQUEST FOR FILING APPLICATION

Under Rule 53(a), (b) & (f)

(No Filing Fee or Oath/Declaration)

(Do NOT use for Provisional or PCT Applications)Use for Design or Utility ApplicationsPATENT
APPLICATION

jc772 U.S. PTO



09/20/00

RULE 53(f) NO DECLARATIONAssistant Commissioner of Patents
and Trademarks
Washington, DC 20231Atty. Dkt. PM 268411
M#

Client Ref.

Date: September 20, 2000

Sir:

1. This is a Request for filing a new Patent Application(☐ Design ☒ Utility) entitled:2. (Complete) Title: Method for the therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and fallopian tube obstruction
without a filing fee or Oath/Declaration but for which is enclosed the following:3. ☒ Abstract 1 page(s).4. 23 Pages of Specification (only spec. and claims); 5. ☐ Specification in non-English language6. 27 Numbered claim(s); and7. ☒ Drawings: 2 sheet(s) ☐ 1 set informal; 8. ☒ formal of size: ☒ A4 ☐ 11"9. **DOMESTIC/INTERNATIONAL** priority is claimed under 35 USC 119(e)/120/365(c) based on the following provisional, nonprovisional and/or PCT international application(s):

Application No.	Filing Date	Application No.	Filing Date
(1) <u>60/155,478</u>	<u>September 23, 1999</u>	(2)	
(3) <u>264771</u>		(4)	
(5)		(6)	

10. **FOREIGN** priority is claimed under 35 USC 119(a)-(d)/365(b) based on filing in _____

Application No.	Filing Date	Application No.	Filing Date
(1)		(2)	
(3)		(4)	
(5)		(6)	

11. _____ (No.) Certified copy (copies): ☐ attached; ☐ previously filed (date) _____
in U.S. Application No. / filed on _____12. ☐ This is a reissue of Patent No. _____13. ☐ See top first page re prior Provisional, National, International application(s) (X box only if info is there and do not complete corresponding item 14 or 15.)14. ☐ **Amend the specification** by inserting before the first line -- This is a ☐ Continuation-in-Part
☐ Divisional ☐ Continuation ☐ Substitute Application (MPEP 201.09) of:14(a) ☐ National Appln. No. / filed --- (M#)14(b) ☐ International Appln. No. PCT/ filed _____ which designated the U.S. --15. ☐ **Amend the specification** by inserting before the first line: --This application claims the benefit of U.S. Provisional Application No. 60/ _____, filed _____ --16. Extension to date: ☐ concurrently filed ☐ not needed ☐ previously filed

17. ☐ Prior application is assigned to

by Assignment recorded _____ Reel _____ Frame _____

18. ☐ Attached:

19. This application is made by the following named inventor(s)

(Double check instructions for accuracy.):

(1) Inventor	Hilde		Riethmuller-Winzen
	First	Middle Initial	Family Name
Residence	Frankfurt	Germany	Germany
	City	State/Foreign Country	Country of Citizenship
Post Office Address			
(include Zip Code)			

(2) Inventor	Jurgen		Engel
	First	Middle Initial	Family Name
Residence	Alzenau	Germany	Germany
	City	State/Foreign Country	Country of Citizenship
Post Office Address			
(include Zip Code)			

(3) Inventor	Ricardo		Felberbaum
	First	Middle Initial	Family Name
Residence	Lubeck	Germany	Germany
	City	State/Foreign Country	Country of Citizenship
Post Office Address			
(include Zip Code)			

(4) Inventor			
	First	Middle Initial	Family Name
Residence			
	City	State/Foreign Country	Country of Citizenship
Post Office Address			
(include Zip Code)			

(5) Inventor			
	First	Middle Initial	Family Name
Residence			
	City	State/Foreign Country	Country of Citizenship
Post Office Address			
(include Zip Code)			

20. NOTE: FOR ADDITIONAL INVENTORS, check box ☐
and attach sheet with same information regarding additional inventors.

Pillsbury Madison & Sutro LLP
Intellectual Property Group

1100 New York Avenue, NW.
Ninth Floor
Washington, DC 20005-3918
Tel: (202) 861-3000
Atty/Sec: ASH/maf

By: Atty: Ann S. Hobbs

Reg. No. 36830

Sig: 

Fax: (202) 822-0944
Tel: (202) 861-3063

NOTE: File in duplicate with 2 post card receipts (PAT-103) & attachments

Method for the therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and fallopian tube obstruction

5

Field of Invention

Endometriosis is one of the most frequently encountered pathologies diagnosed amongst gynecological patients. For example, between 10% and 25% of women presenting with gynecological symptoms in UK and in the USA are affected. Clinical diagnosis is made usually by laparoscopic observation of hemorrhagic or fibrotic foci on the pelvic organs. The ectopic endometrial tissue responds to ovarian hormones undergoing cyclic changes. The cyclical bleeding from the endometric deposit contributes to a local inflammatory reaction. Endometriosis commonly affects women during their childbearing years with an incidence of at least 1% (see Shaw, R.W. (1993), *An Atlas of Endometriosis*. The Parthenon Publishing Group).

Endometriosis is usually classified into endometriosis (genitalis) interna (adenomyosis), endometriosis genitalis externa and endometriosis extragenitalis.

20 Chronic pelvic pain may occur either in relation to endometriosis or as an independent disease.

Fallopian tube obstruction (FTO) is a relatively common disease and may account to
25 for up to 20 % of cases of tubal infertility (see Winfield, A.C. et al., Apparent cornual
occlusion in hysterosalpingography: Reversal by glucagon. AJR Am J Roentgenol
1982; 139: 525 – 527).

Background information and Prior Art

30

Sampson suggested that menstrual regurgitation and subsequent implantation of endometrial tissue on the peritoneal face results in endometriosis [Sampson, J.A.

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-2-

(1927), Peritoneal endometriosis due to menstrual dissemination of the endometrial tissue into the peritoneal cavity. *Am. J. Obstet. Gynecol.*, 14, 422.]

Several aetiologic factors may be involved in the pathogenesis of endometriosis :

5 Dmowski et. al. suggested that genetic and immunological factors lead to endometriosis [Dmowski, W.P., Steele, R.W. and Baker, G.F. (1981). Deficient cellular immunity in endometriosis. Am. J. Obstet. Gynecol., 141, 377]

Vascular and lymphatic embolization to distant sites has been demonstrated and explains the (rare) finding of endometriosis outside the peritoneal cavity, e.g. skin, lung, kidney.

Cells lining the Müllerian duct arise from primitive cells which differentiate into peritoneal cells and the cells on the surface of the ovaries. It is proposed that these adult cells undergo de-differentiation back to their primitive origin and then transform into endometrial cells [Levander, G. (1941), Bone formation by induction. An experimental study. Arch. Klin. Chir., 202, 497]

Dysmenorrhea, acute or chronic pelvic pain, dyspareunia, and infertility perform the most frequent clinical symptoms reported.

FTO represents a heterogenous group of underlying pathology, preliminary intrinsic
occlusion or extrinsic compression from estrogen-sensitive disorders, such as
endometriosis, adenomyosis, endosalpingiosis, and myomata. FTO is frequently
diagnosed by hysterosalpingography, besides laparoscopy.

First choice of treatment comprises laparoscopic removal of endometric lesions. This procedure may be followed by the treatment with Danazol or LHRH agonist (for a period of six months). Women being treated with Danazol might experience gastrointestinal and hepatic disorders as well as severe androgenic side effects.

It was also proposed from a theoretical viewpoint for treatment of endometriosis and uterine myoma to use the immediate suppression by administration of a LHRH antagonist to reducing the duration of treatment and faster improvement of subjective symptoms [Th. Reissmann et al. Human Reproduction vol. 10 No. 8 pp.1974-1981,(1995)]

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-3-

Further Hodgen teaches in the US Patent 5,658,884 a regime for therapeutic management of a gonadal dependent condition by reducing the estrogen supply by means of long-term administration of an GnRH antagonist for 6 months or longer in an amount effective to inhibit proliferation of endometrial tissue without substantially stopping the production of endogenous estrogen. For this purpose, Hodgen teaches such a regimen or dose of GnRH antagonist to achieve a 24 hour serum estradiol level in the range of about 25 to 50 and preferably about 35 to 45 pg/ml. However, Hodgen does not describe estradiol serum levels oscillating between 50 pg/ml and 75 pg/ml. Moreover, Hodgen only teaches in the US Patent 5,658,884 a continuous long-term treatment (on a daily or periodic basis, the latter meaning a weekly or monthly administration) but not a short-term induction treatment for only 4 to 12 weeks. Hodgen also does not describe any combination therapy comprising the GnRH antagonist in the treatment of endometriosis. The treatment is only described on monkeys and also includes the performance of a costly and repeated progesterone challenge test to provide an 24 hour average serum estradiol level of 30 to 50 pg/ml.

As a consequence of the flare-up effect of LHRH-agonistic therapy an exacerbation of symptoms might occur during some days. Following prolonged treatment which is required to avoid the re-proliferation of endometric tissue hormonal withdrawal symptoms as well as demineralization of bones occur.

Therefore, effective drug therapy should immediately reduce the residual extrauterine endometrial tissue present after laparoscopic surgery. Duration of therapy should be only 4 to 12 weeks without the occurrence of any major hormonal withdrawal symptoms or ovarian cyst formation.

LHRH antagonists exert an immediate onset of hormonal suppression, and therefore benign gynecological tumors, such as uterine fibroids decrease within short time [Human Reproduction 1998, 13]

Object of the Invention

The present invention relates to the improvement of the medical treatment of extrauterine proliferation of endometrial tissue, i. e. the administration of LHRH antagonists in patients with clinical symptoms of endometriosis, the improvement consisting of :

- 5 immediate reduction of ectopic endometrial tissue
 immediate cessation of symptoms, e.g. severe pain, chronic pelvic pain and
 dysmenorrhea
 prevention of any progress of the disease
 avoidance of hormonal withdrawal symptoms
- 10 prevention of ovarian cyst formation, demineralization of bones as well as of
 gastrointestinal or hepatic disorders.

The inventive medical therapy can start in the early to mid follicular phase, preferably on cycle day one to three. During the treatment the estradiol serum concentration levels are kept between 35 pg/ml and 80 pg/ml, preferably between about 45-75 pg/ml, more preferably between about 50-75 pg/ml. The LHRH antagonist is administered only for 4 to 12 weeks (short-term induction treatment), either by daily, weekly or monthly administration. Following the short-term induction treatment, the administration of a contraceptive, a non-steroidal anti-rheumatic, an analgetic, an androgen other than 17-alpha-alkyl substituted testosterone or any combinations thereof is provided according to the present invention.

Summary of the Invention

- 25 In the treatment of extrauterine endometrial tissue with an LHRH antagonist, therapy is started on menstrual cycle day one to three. Before starting LHRH-antagonist therapy the diagnosis is performed by laparoscopy. In cases of severe pain, LHRH antagonist therapy might be initiated without prior laparoscopy.

Therapy will continue until clinical symptomatology has resolved and no proliferation of the endometrium is seen. Due to the immediate onset of suppression of the gonadotropins LH and FSH as well of sex steroids estradiol and progesterone no

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-5-

further proliferation of the endometrium occurs. Benign tumors or other sex steroid dependent lesions, like endometriosis decrease within four to twelve weeks of therapy. Due to the lack of flare-up no ovarian cysts develop.

- 5 Furthermore, no hormonal withdrawal symptoms are seen as the estradiol values are kept in the range of the early follicular phase of 35 to 80 pg/ml, preferably between about 45-75 pg/ml, more preferably between about 50-75 pg/ml without further increase or decrease. No titering of the dosage of the LHRH antagonist, e.g. by conducting a costly progesterone challenge test, is necessary.

10

The method of therapeutic management of extrauterine proliferation of endometrial tissue the improvement according to the invention therefore embraces:

immediate reduction of ectopic endometrial tissue

prevention of any progress of the disease

- 15 avoidance of hormonal withdrawal symptoms

prevention of ovarian cyst formation, demineralization of bones as well as of gastrointestinal or hepatic disorders

- 20 start of medical therapy on cycle day one to three and maintenance of estradiol levels at values of the early follicular phase throughout the entire duration of treatment by means of administration of a LHRH antagonist wherein the antagonist is preferably cetrorelix, leuprorelin, goserelin, antide or abarelix. The antagonist can also be the LHRH antagonist D-63153 (Ac-D-Nal-D-pCl-Phe-D-Pal-Ser-N-Me-Tyr-D-Hci-Nle-Arg-Pro-D-Ala-NH₂) as described in the German Patent Application No. 199 11 771.3 filed on March 11, 1999.

25

The LHRH antagonist may be given for 4 to 12 weeks in a weekly dose of 3 to 10 mg per week or for 4 to 12 weeks in a daily dose of 0.25 mg to 0.5 mg/day.

- 30 It is also possible to give the LHRH antagonist 4 to 12 weeks in a monthly dose of 12 to 40 mg per month.

In a repeat therapeutic treatment the LHRH antagonist is given for 4 to 12 weeks and the treatment is repeated two or three times a year, whereby a repeated

000350" 9479950

treatment does not follow directly after a short-term induction treatment. Usually a period of time of weeks or months, where no LHRH antagonist is administered, is between the end of the short-term induction treatment and the start of the repeat treatment.

5

To demonstrate the feasibility to maintain a low estradiol secretion under adjusted LHRH- antagonist treatment so that a therapeutic suppression occurs without withdrawal symptoms nine patients with confirmed endometriosis were treated with 3 mg of Cetrorelix acetate s.c. by weekly administration for 8 weeks. While patients compliance was excellent avoiding any hot flushes or other withdrawal symptoms and without any progress of the disease confirmed by 2nd look laparoscopy the mean estradiol serum concentrations oscillated between 37 pg/ml and 64 pg/ml, preferably between 45-75 pg/ml, more preferably between about 50-75 pg/ml. Histological biopsies showed no proliferation of the endometrium at the end of treatment. No ovarian cyst formation occurred.

The figure 1 shows the continuous estradiol suppression to values of the early follicular phase (range of 35 pg/ml to 80 pg/ml, preferably between 45-75 pg/ml, more preferably between about 50-75 pg/ml) obtained in patients with endometriosis by a weekly dose of 3 mg of Cetrorelix (LHRH antagonist) for 8 weeks. Immediate and continuous suppression of estradiol levels is obtained without any signs of estradiol withdrawal symptoms and without proliferation of the endometrium at the end of treatment.

25 Fig. 2 shows estradiol serum levels after administration of cetrorelix at a weekly dose of 1 mg resp. 3 mg once per week. The estradiol serum levels are between about 35-80 pg/ml, preferably between about 45-75 pg/ml, more preferably between about 50-75 pg/ml.

30 The endometriosis patient with distinctive symptomatic pain is suffering from a chronic disease. Surgical methods in sense of curative therapy as well as medicinal treatment to suppress the sexual steroid secretion of the patient often result only in a

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-7-

temporary improvement. The relapse rate of the discomforts is very high and about 70% within 5 years after finishing therapy (Schweppe, 1999).

At the same time the radical surgical therapy and the suppression of the estrogen secretion leads to considerable side effects. The radical surgical therapy in sense of hysterectomy with bilateral adnexectomy is no adequate therapy for the younger, premenopausal woman. The chronical lack of estrogen leads to the following vegetative symptoms: hot flashes, sweating, dryness of the vagina, depressive feelings and also holds the risk of osteoporosis. The alternative therapy with the synthetic steroidal compound Danazol may cause virilizing symptoms because of the androgenic effect.

Aim of the medicinal therapy of patients with endometriosis with symptomatic pain is to obtain a treatment without side effects, especially avoiding the negative effects of estrogen suppression and which is long-lasting after finishing therapy. The specific pharmacological mode of action of LHRH antagonists allows new possibilities for treatment of endometriosis.

The weekly administration of an adequate dose of an LHRH antagonist, e.g. 3 mg Cetrotide® s.c./ per week over a period of eight weeks leads to a controlled suppression of estrogen secretion so that serum concentrations between about 35 pg/ml and about 80 pg/ml, preferably between about 45-75 pg/ml, more preferably between about 50-75 pg/ml are obtained. In this serum concentration range no vegetative symptoms arise. Also the development of osteoporosis can be avoided. The symptomatic pain will be effectively suppressed in all stages of the disease (rAFS I – IV). In the stages rAFS I – II a clinical regression of the disease in sense of decrease of the implantation area is noticed (Felberbaum et. al., 2000).

In a preferred embodiment of this invention, after this treatment period of eight to twelve weeks the patient could take a contraceptive, preferably an oral contraceptive, preferably with gestagen components, unless there is a wish for pregnancy. In this connection combinations with Lynestronol 2 mg with 0,04 mg

000000 " 9479360

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-8-

Ethinylestradiol or 2,5 mg Lynestrenol with 0,05 mg of Ethinylestradiol (e.g. Yermonil®, Lyn-ratiopharm-Sequenz®) have to be mentioned.

A combination therapy with androgens other than 17-alpha-alkyl substituted
5 testosterone such as danazol may also be applied subsequently to the short-term
induction regimen with the LHRH antagonist either alone or in combination with non-
steroidal anti-rheumatics and/or analgetics. An example for a suitable androgene is
halotestin™ (fluoximesterone).

10 The treatment with a contraceptive, preferably an oral contraceptive, preferably
containing gestagens, should be individually continued until typical pain sensation
occurs. In this stage the patient will have relatively small menstrual bleeding as an
effect of the gestagen component of this contraceptive, preferably oral contraceptive.
For covering also the especially critical pre-menstrual and menstrual days with
15 regard to pain sensation in this phase a concomitant medication with appropriate
non-steroidal anti-rheumatic drugs, e.g. diclophenac, ibuprofen, indometacin,
oxicam derivates or acetylsalicylic acid may be given. Also an analgetic such as
flupirtinmaleat (Katadolon®) can be administered.

20 If further pain symptoms occur during this combination therapy with gestagenic
contraceptives, preferably oral contraceptives, a daily, weekly or monthly therapy
with the adequate dose of an LHRH antagonist as described above may be
repeated. Detailed information on the respective treatment options are given below.
If the patient is absolutely free of pain treatment can be changed to gestagenic
25 contraceptive, preferably oral contraceptive,s in combination with concomitant
medication of appropriate non-steroid anti-rheumatic drugs or analgetics.

This therapy using the intermittent administration of an LHRH antagonist leads to a
new and innovative unlimited treatment without side effects and lowers treatment
30 burden for the patient significantly.

0966146-092000

Pharmaceutical Formulations Suitable for Treatment

Pharmaceutical formulations of the LHRH antagonist suitable for the therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and fallopian tube obstruction may be for example

a) acetate salt formulations in the concentration of 1 mg/1 ml or lower where the powder may be dissolved in Water for Injection (Wfi) or in Gluconic Acid (GA);

b) acetate salt formulations in the concentration of 1.5 mg/1 ml to 5.0 mg /1 ml, preferably 2.5 mg/1 ml where the powder may be dissolved in Water for Injection (Wfi) or in Gluconic Acid (GA);

c) pamoate salt formulations in the concentration of 10 mg/1 ml to 30 mg/1 ml, preferably 15 mg/1 ml where the lyophilisate powder may be dissolved in Gluconic Acid (GA) or in Water for Injection (Wfi).

According to one aspect of the present invention in the method of therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO), the improvement consisting of administration of an LHRH antagonist in the form of a short-term induction treatment for a period of about 4 to 12 weeks to a patient in need of such treatment, whereby subsequently the administration of the LHRH antagonist is ceased, is provided.

The duration of the short term induction treatment is about 4 to about 12 weeks, that means that the treatment can be between about 28 to about 84 days or from about one to about three months.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, wherein the LHRH antagonist is administered such that the estrogen serum concentration level is between about 35 pg/ml and

-10-

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the short-term induction treatment with the LHRH antagonist is followed by administration of a contraceptive, preferably an oral contraceptive.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the short-term induction treatment with the LHRH antagonist is followed by administration of a non-steroidal anti-rheumatic agent.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the short-term induction treatment with the LHRH antagonist is followed by administration of an analgetic.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the short-term induction treatment with the LHRH antagonist is followed by administration of an androgen other than a 17-alpha-alkyl substituted testosterone.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the short-term induction treatment with the LHRH antagonist is followed by the combined or separate administration of one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof.

According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-11-

administered starting in the early to mid follicular phase, preferably on cycle day one to three.

5 According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is selected from the group consisting of cetrorelix, teverelix, ganirelix, antide, abarelix and D-63153.

10 According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a weekly dose of 3 to 10 mg per week.

15 According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a daily dose of 0.25 mg to 0.5 mg/day.

20 According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a monthly dose of 12 to 40 mg per month.

25 According to another aspect of the present invention in a method as mentioned above the improvement is provided, characterized in that the LHRH antagonist is given for the induction treatment during 4 to 12 weeks and the treatment is repeated two or three times a year.

30 According to a further aspect of the present invention a pharmaceutical composition for treating extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO) comprising an LHRH antagonist and optionally one or more agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic agent, an

0966145-092000

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-12-

androgen agent other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, optionally together with pharmaceutically acceptable excipients, whereby the LH-RH antagonist is administered to a patient in need thereof in a short term induction treatment for a period of about 4 to 12 weeks, then the administration of the LH-RH antagonist is ceased and optionally the one or more agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, are administered together or separately to the patient is provided.

Suitable excipients and dosage forms are for example described by K.H. Bauer, K.-H. Frömming and C. Führer, Lehrbuch der Pharmazeutischen Technologie, 6th edition, Stuttgart 1999, pages 163-186 (excipients) and pages 227-386 (dosage forms), including the references as cited therein.

The LH-RH antagonist can be administered for example subcutaneous (s.c.), intramuscular (i.m.) or inhalative. The agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof can be administered as known in the art (see for example the German, European or U.S. pharmacopoeia), preferably oral or inhalative.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is administered such that the estrogen serum concentration level is between about 35 pg/ml and about 80 pg/ml, preferably between about 45–75 pg/ml, more preferably about 50–75 pg/ml.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a contraceptive, preferably an oral contraceptive.

Applicant: ASTA Medica AG

Applicant's ref. no.: 99/16 PH/Ba

-13-

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a non-steroidal anti-rheumatic agent.

5

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an analgetic.

10 According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an androgen other than a 17-alpha-alkyl substituted testosterone.

15 According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the short-term induction treatment with the LHRH antagonist is followed by the combined or separate administration of one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an
20 androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is administered starting in the early to mid follicular phase, preferably on cycle day one to three.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is selected from the group consisting of cetrorelix, teverelix, ganirelix, antide, abarelix and D-63153.

30

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is administered during

the short-term induction treatment for 4 to 12 weeks at a weekly dose of 3 to 10 mg per week.

According to another aspect of the present invention, a pharmaceutical composition
5 as mentioned above is provided wherein the LHRH antagonist is administered during
the short-term induction treatment for 4 to 12 weeks at a daily dose of 0.25 mg to 0.5
mg/day.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a monthly dose of 12 to 40 mg per month.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided wherein the LHRH antagonist is given for the induction treatment during 4 to 12 weeks and the treatment is repeated two or three times a year.

According to another aspect of the present invention, a pharmaceutical composition as mentioned above is provided, wherein the the one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, are in the same or separate dosage forms.

25 According to another aspect of the present invention, a use of an LH-RH antagonist for the preparation of a medicament for the therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO), whereby the LHRH antagonist is administered in the form of a

30 short-term induction treatment for a period of about 4 to 12 weeks to a patient in need of such treatment and then the administration of the LHRH antagonist is ceased, is provided.

-15-

5

10

15

20

25

30

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-16-

According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is administered starting in the early to mid follicular phase, preferably on cycle day one to three.

- 5 According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is selected from the group consisting of cetrorelix, teverelix, ganirelix, antide, abarelix and D-63153.

- 10 According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a weekly dose of 3 to 10 mg per week.

- 15 According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a daily dose of 0.25 mg to 0.5 mg/day.

- 20 According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a monthly dose of 12 to 40 mg per month.

- 25 According to another aspect of the present invention, a use of an LH-RH antagonist as mentioned above is provided, wherein the LHRH antagonist is given for the induction treatment during 4 to 12 weeks and the treatment is repeated two or three times a year.

- 30 According to another aspect of the present invention, a use of an LH-RH antagonist and one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone, or any combinations thereof, for the preparation of a medicament for the therapeutic

0966145-090000

management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO), whereby the LHRH antagonist is administered in the form of a short-term induction treatment for a period of about 4 to 12 weeks to a patient in need of such treatment, then the administration of the LHRH antagonist is ceased and the one or more active agent selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone, or any combinations thereof, are administered together or separately to the patient, is provided.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is administered such that the estrogen serum concentration level is between about 35 pg/ml and about 80 pg/ml, preferably between about 45-75 pg/ml, more preferably about 50-75 pg/ml.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a contraceptive, preferably an oral contraceptive.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a non-steroidal anti-rheumatic agent.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an analgetic.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the short-

term induction treatment with the LHRH antagonist is followed by administration of an androgen other than a 17-alpha-alkyl substituted testosterone.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the short-term induction treatment with the LHRH antagonist is followed by the combined or separate administration of one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is administered starting in the early to mid follicular phase, preferably on cycle day one to three.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is selected from the group consisting of cetrorelix, teverelix, ganirelix, antide, abarelix and D-63153.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a weekly dose of 3 to 10 mg per week.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a daily dose of 0.25 mg to 0.5 mg/day.

According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-19-

antagonist is administered during the short-term induction treatment for 4 to 12 weeks at a monthly dose of 12 to 40 mg per month.

- 5 According to another aspect of the present invention the use of an LH-RH antagonist and one or more active agents as mentioned above is provided, wherein the LHRH antagonist is given for the induction treatment during 4 to 12 weeks and the treatment is repeated two or three times a year.

000260-94799960

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba

-20-

Claims :

1. In the method of therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO),
5 the improvement consisting of administration of an LHRH antagonist in the form of a short-term induction treatment for a period of about 4 to 12 weeks to a patient in need of such treatment, whereby subsequently the administration of the LHRH antagonist is ceased.
- 10 2. A method according to claim 1 wherein the LHRH antagonist is administered such that the estrogen serum concentration level is between about 35 pg/ml and about 80 pg/ml, preferably between about 45-75 pg/ml, more preferably about 50-75 pg/ml.
- 15 3. A method according to claim 1 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a contraceptive, preferably an oral contraceptive.
- 20 4. A method according to claim 1 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a non-steroidal anti-rheumatic agent.
- 25 5. A method according to claim 1 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an analgetic.
6. A method according to claim 1 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an androgen other than a 17-alpha-alkyl substituted testosterone.
- 30 7. A method according to claim 1 wherein the short-term induction treatment with the LHRH antagonist is followed by the combined or separate administration of one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an

0966146-093000

Applicant: AS I A Medica AG

Applicant's ref. no.: 99/16'PH/Ba

-21-

analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof.

8. A method according to claim 1 wherein the LHRH antagonist is administered starting in the early to mid follicular phase, preferably on cycle day one to three.
9. A method according to claim 1 wherein the LHRH antagonist is selected from the group consisting of cetrorelix, toverelix, ganirelix, antide, abarelix and D-63153.
10. A method according to claim 1 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a weekly dose of about 3 to 10 mg per week.
11. A method according to claim 1 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a daily dose of about 0.25 mg to 0.5 mg/day.
12. A method according to claim 1 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a monthly dose of about 12 to 40 mg per month.
13. A method according to claim 1 wherein the LHRH antagonist is given for the induction treatment during about 4 to 12 weeks and the treatment is repeated two or three times a year.
14. A pharmaceutical composition for treating extrauterine proliferation of endometrial tissue, chronic pelvic pain and/or fallopian tube obstruction (FTO) comprising an LHRH antagonist and optionally one or more agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, optionally together with pharmaceutically acceptable excipients, whereby the LH-RH antagonist is administered to a patient in need thereof in a short term induction treatment for a

period of about 4 to 12 weeks, then the administration of the LH-RH antagonist is ceased and optionally the one or more agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, are administered together or separately to the patient.

15. Pharmaceutical composition according to claim 14 wherein the LHRH antagonist is administered such that the estrogen serum concentration level is between about 35 pg/ml and about 80 pg/ml, preferably between about 45–75 pg/ml, more preferably about 50–75 pg/ml.
16. Pharmaceutical composition according to claim 14 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a contraceptive, preferably an oral contraceptive.
17. Pharmaceutical composition according to claim 14 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of a non-steroidal anti-rheumatic agent.
18. Pharmaceutical composition according to claim 14 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an analgetic.
19. Pharmaceutical composition according to claim 14 wherein the short-term induction treatment with the LHRH antagonist is followed by administration of an androgen other than a 17-alpha-alkyl substituted testosterone.
20. Pharmaceutical composition according to claim 14 wherein the short-term induction treatment with the LHRH antagonist is followed by the combined or separate administration of one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16' PH/Ba

-23-

anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof.

21. Pharmaceutical composition according to claim 14 wherein the LHRH antagonist is administered starting in the early to mid follicular phase, preferably on cycle day one to three.
22. A pharmaceutical composition according to claim 14 wherein the LHRH antagonist is selected from the group consisting of cetrorelix, teverelix, ganirelix, antide, abarelix and D-63153.
23. Pharmaceutical composition according to claim 14 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a weekly dose of about 3 to about 10 mg per week.
24. A pharmaceutical composition according to claim 14 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a daily dose of about 0.25 mg to about 0.5 mg/day.
25. Pharmaceutical composition according to claim 14 wherein the LHRH antagonist is administered during the short-term induction treatment for about 4 to 12 weeks at a monthly dose of about 12 to 40 mg per month.
26. Pharmaceutical composition according to claim 14 wherein the LHRH antagonist is given for the induction treatment during about 4 to 12 weeks and the treatment is repeated two or three times a year.
27. Pharmaceutical composition according to claim 14, wherein the the one or more active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof, are in the same or separate dosage forms.

Applicant: ASTA Medica AG
Applicant's ref. no.: 99/16 PH/Ba


-24-

Abstract

- 5 The present invention provides a method for therapeutic management of extrauterine proliferation of endometrial tissue, chronic pelvic pain and fallopian tube obstruction by short term induction treatment with an LH-RH antagonist for 4 to 12 weeks. According to another aspect of the present invention, the short term LH-RH treatment is followed by the combined or separate administration of one or more
10 active agents selected from the group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof. According to a further aspect of the present invention a pharmaceutical composition comprising an LHRH antagonist and one or more active agents selected from the
15 group consisting of a contraceptive, preferably an oral contraceptive, a non-steroidal anti-rheumatic agent, an analgetic, an androgen other than a 17-alpha-alkyl substituted testosterone or any combinations thereof are provided.

figure 1

CTT in Endometriosis
estradiol [pg/ml]



Celorelix in patients with endometriosis 3 mg celorelix per week starting on cycle day one to three
Estradiol serum levels during 8 weeks therapy

Cetrorelix 1mg and 3 mg once per week Time Courses of Serum E₂ Levels (Median Values)

